

21st Century Topical Management of Superficial Pyoderma



Galia Sheinberg, DVM, DLACVD

Dr. Sheinberg is a board certified veterinary dermatologist who works in a busy referral practice in Mexico City, Centro Veterinario Mexico. She consults on 240 to 260 dermatology cases per month. And, successfully manages over 90% of her superficial pyoderma cases using topical therapy in lieu of antibiotics.



Antibiotic bacterial resistance is a growing problem and poses a therapeutic challenge in human and veterinary medicine. Antibiotics are used not only to treat bacterial infections in human patients and animals but also the food industry. Animals and plants are treated with antimicrobials to prevent and treat disease and improve productivity. Resistance is not a new phenomenon. It has always been a survival strategy for bacteria, but antibiotics create selective pressure that promotes and induces rapid bacterial resistance. Some bacteria have become resistant to many classes of antibiotics; such is the case with *Staphylococcus pseudintermedius*, which is the most common bacteria found in canine pyoderma. Antibiotics that were commonly used to treat dogs with skin infections are often no longer effective, and the treatment of these infections has also become more complicated and expensive, requiring bacterial cultures with antibiotic sensitivities.¹

Unfortunately, the antibiotics that are used for treating multi-resistant bacteria are either toxic, expensive, promote more resistance, or are complicated to use, posing a big challenge for veterinarians. Furthermore, empirical drug selection for systemic therapy is always contraindicated when an MRS infection is suspected.²

Current recommendations and guidelines for treating superficial and surface pyoderma in dogs are to use topical therapy first. Patients may be treated locally using wipes, sprays, ointments, and mousses or bathed for generalized problems. Several studies have proven that the use of topical chlorhexidine or a combination of chlorhexidine and miconazole can be as or more effective than the use of antibiotics. Consideration of client compliance is paramount when choosing topical therapy.^{2,3,4,5}

When faced with a patient with pyoderma, veterinarians need to choose carefully the best strategy for the management of that particular patient. For example, how involved is the owner, how much money are they willing to spend on treatment, how much time do they have available, and how concerned is the client with their pet's disease? Client education is not only essential to achieving proper treatment and compliance for patients, but also has a substantial effect on disease prevention management and patient welfare. Generally, clients prefer fast results and less work making antibiotics more attractive. Taking the time to explain why topical therapy is the better option also requires time and patience, but can yield great results.

Treating bacterial infection with topical therapy requires management of concurrent pruritus and inflammation using an appropriate anti-inflammatory and anti-itch product. Recent publications suggest using steroids initially when inflammation is more severe and transitioning to more targeted therapies such as oclacitinib or Lokivetmab that have no or fewer side effects.⁶ Choosing what to use requires a tailored and individualized therapy; some patients are not able to use steroids because of concurrent disease. Oclacitinib has been demonstrated to work as fast or faster than steroids in a previous study.⁷

In recent years in our practice, we have managed hundreds of patients using primarily topical therapy with excellent results avoiding unnecessary use of antibiotics. Generalized pyodermas respond very well to shampoo therapy. Deciding how often to bathe depends on the type of coat and size of the dog. Severe cases might benefit from bathing every other day, however this is not always possible. Using sprays or mousse on large areas are an easy way of medicating those areas without bathing.

REFERENCES:

1. Schwartz, Anette Loeffler, Kristina Kadlec. Bacterial resistance to antimicrobial agents and their impact on veterinary and human medicine. *Vet Dermatol* 2017; 28: 82–e19
2. Daniel O. Morris, Anette Loeffler, Meghan F, et al. Recommendations for approaches to methicillin-resistant staphylococcal infections of small animals: diagnosis, therapeutic considerations, and preventative measures. *Clinical Consensus Guidelines of the World Association for Veterinary Dermatology*. *Vet Dermatol* 2017; 28: 304–e69
3. Stefano Borio, Silvia Colombo, Giuseppe La Rosa, et al. Effectiveness of a combined (4% chlorhexidine digluconate shampoo and solution) protocol in MRS and non-MRS canine superficial pyoderma: a randomized, blinded, antibiotic-controlled study. *Vet Dermatol* 2015; 26: 339–e72
4. Loeffler A, Cobb MA, Bond R. Comparison of a chlorhexidine and a benzoyl peroxide shampoo as sole treatment in canine superficial pyoderma. *Vet Rec* 2011; 169: 249.
5. Murayama N, Nagata M, Terada Y, et al. Comparison of two formulations of chlorhexidine for treating canine superficial pyoderma. *Vet Rec* 2010; 167: 532–533.
6. Thierry Olivry, Frane Banovic. Editorial treatment of canine atopic dermatitis: time to revise our strategy. *Vet Dermatol* 2019; 30: 87–90
7. Andrea J. Gonzáles, Thimoty J. Fleck, William R. Humphrey, et al. IL-31-induced pruritus in dogs: a Novel experimental model to evaluate anti-pruritic effects of canine therapeutics. *Vet Dermatol* 2016; 27-34 E 10